

Curriculum Vitae

Academic Work

2004-2022/Jan

Personal Information

Name: Toshiro Inaba, M.D. Ph.D.

Place of Birth: Kumamoto, Japan

Degrees:

M.D. 2004 (University of Tokyo, Faculty of Medicine)

Ph.D. 2014 (University of Tokyo, Internal Medicine)

Work Experience

2020- Karuizawa Municipal Hospital, Department of General medicine (Nagano, JAPAN)

2021- Assistant director of Karuizawa hospital

- 2020- University of Shinshu, Associate Professor of the Shinshu University Research Center for Social Systems (Nagano JAPAN)
- 2020-Tohoku University of Art and Design, Visiting Professor and Yamagata Biennale 2020 Art Director
- 2020-UTokyoRcast(Research Center for Advanced Science and Technology, The University of Tokyo), Visiting Researcher

2009 -2020 University of Tokyo, Department of Cardiovascular medicine (Tokyo, JAPAN)

2014- Assistant Professor, Division of Cardiology, The Tokyo University Hospital (Tokyo, JAPAN)

2006-2009 The Cardiovascular Institute Hospital (Tokyo, JAPAN)

2004-2006 Aizawa hospital (Nagano, JAPAN)

Education

2010-2014 University of Tokyo School of Medicine (Doctor of Medicine)

1998-2004 Graduated School of Medicine and Faculty of Medicine, The University of Tokyo (Doctor of Philosophy)

Honors and Awards

2012 AHA (Los Angeles), Cardiopulmonary, Critical Care, Perioperative and Resuscitation (3CPR) Best Abstract Award.

Inaba T, Yao A, Hatano M, Maki H, Imamura T, Kinugawa K, Komuro I. Hemodynamic Characteristics in Systemic Sclerosis Patients with Various Stages of Pulmonary Arterial Hypertension.

Publication

Inaba T, Yao A, Nakao T, Hatano M, Maki H, Imamura T, Shiga T, Yamazaki T, Sonoda M, Kinugawa K, Shiota T, Suzuki J, Takenaka K, Hirata Y, Nagai R. Volumetric and functional assessment of ventricles in pulmonary hypertension on 3-dimensional echocardiography. *Circ J*. 2012;77(1):198-206.

Peer-Reviewed Manuscripts

Clinical Outcome of Closure of a Small Atrial Septal Defect in a Patient with Pulmonary Arterial Hypertension.

Umei M, Saito A, Soma K, **Inaba T**, Uehara M, Komuro I, Yao A.
Intern Med. 2021 Aug 31.

Association of serum CCL20 levels with pulmonary vascular involvement and primary biliary cholangitis in patients with systemic sclerosis.

Ikawa T, Miyagawa T, Fukui Y, Minatsuki S, Maki H, **Inaba T**, Hatano M, Toyama S, Omatsu J, Awaji K, Norimatsu Y, Watanabe Y, Yoshizaki A, Sato S, Asano Y.
Int J Rheum Dis. 2021 May;24(5):711-718.

Factors associated with left ventricular reverse remodelling after percutaneous coronary intervention in patients with left ventricular systolic dysfunction.

Adachi Y, Kiyosue A, Ando J, Kawahara T, Kodera S, Minatsuki S, Kikuchi H, **Inaba T**, Kiriyaama H, Hirose K, Shinohara H, Saito A, Fujiwara T, Hara H, Ueda K, Sakakura K, Hatano M, Harada M, Takimoto E, Akazawa H, Morita H, Momomura SI, Fujita H, Komuro I.
Sci Rep. 2021 Jan 8;11(1):239.

Balloon Valvuloplasty to Treat Adult Symptomatic Pulmonary Valve Stenosis with Sequential Follow-Up Using Cardiac Magnetic Resonance Imaging in Combination with Echocardiography.

Ishii S, Fujiwara T, Ando J, Takeda N, **Inaba T**, Inuzuka R, Komuro I.

Int Heart J. 2020 Sep 29;61(5):1075-1078.

The Clinical Efficacy of Endothelin Receptor Antagonists in Patients with Pulmonary Arterial Hypertension.

Maki H, Hara T, Tsuji M, Saito A, Minatsuki S, **Inaba T**, Amiya E, Hosoya Y, Hatano M, Morita H, Yao A, Kinugawa K, Komuro I.

Int Heart J. 2020;61(4):799-805.

Fractional exhaled nitric oxide in adult congenital heart disease.

Saito A, Amiya E, Soma K, **Inaba T**, Maki H, Hatano M, Yao A, Morita H, Komuro I.

Nitric Oxide. 2020 Aug 1;100-101:45-49.

Regular Treatment Strategy with a Large Amount of Carvedilol for Heart Failure Improves Biventricular Systolic Failure in a Patient with Repaired Tetralogy of Fallot.

Soma K, Yao A, Saito A, **Inaba T**, Ishikawa Y, Hirata Y, Komuro I.

Int Heart J. 2018 Sep 26;59(5):1169-1173.

Combined Surgical and Medical Therapy for Candida Prosthetic Endocarditis in a Patient with Repaired Tetralogy of Fallot.

Kubota K, Soma K, Uehara M, **Inaba T**, Saito A, Takeda N, Hatano M, Morita H, Inuzuka R, Hirata Y, Yao A, Komuro I.

Int Heart J. 2018 Jul 31;59(4):877-880.

Diastolic Tricuspid Regurgitation Related to Atrial Systole With a Bileaflet Mechanical Valve.

Saito A, Soma K, Kojima T, **Inaba T**, Yao A, Komuro I.

Circ J. 2018 Oct 25;82(11):2921-2922.

Prevalence of primary Sjögren's syndrome in patients undergoing evaluation for pulmonary arterial hypertension.

Sato T, Hatano M, Iwasaki Y, Maki H, Saito A, Minatsuki S, **Inaba T**, Amiya E, Fujio K, Watanabe M, Yamamoto K, Komuro I.

PLoS One. 2018 May 15;13(5):e0197297.

Serum levels of interleukin-18-binding protein isoform a: Clinical association with inflammation and pulmonary hypertension in systemic sclerosis.

Nakamura K, Asano Y, Taniguchi T, Minatsuki S, **Inaba T**, Maki H, Hatano M, Yamashita T, Saigusa R,

Ichimura Y, Takahashi T, Toyama T, Yoshizaki A, Miyagaki T, Sugaya M, Sato S.
J Dermatol. 2016 Aug;43(8):912-8.

Novel Scoring System to Predict Ineligibility for Bridge to Implantable Left Ventricular Assist Device as Destination Therapy Before Extracorporeal Ventricular Assist Device Implantation - For the Coming Era of Destination Therapy in Japan.

Nitta D, Kinugawa K, Imamura T, Endo M, **Inaba T**, Maki H, Amiya E, Hatano M, Kinoshita O, Nawata K, Kyo S, Ono M.
Circ J. 2016;80(2):387-94.

An Experience of Landiolol Use for an Advanced Heart Failure Patient With Severe Hypotension.

Nitta D, Kinugawa K, Imamura T, Endo M, Amiya E, **Inaba T**, Maki H, Hatano M, Komuro I.
Int Heart J. 2015;56(5):564-7.

Quality of Life and Influential Factors in Patients Implanted With a Left Ventricular Assist Device.

Kato NP, Okada I, Imamura T, Kagami Y, Endo M, Nitta D, Fujino T, Muraoka H, Minatsuki S, Maki H, **Inaba T**, Kinoshita O, Nawata K, Hatano M, Yao A, Kyo S, Ono M, Jaarsma T, Kinugawa K.
Circ J. 2015;79(10):2186-92.

A case of interferon- α -induced pulmonary arterial hypertension after living donor liver transplantation.

Ko T, Hatano M, Nitta D, Muraoka H, Minatsuki S, Imamura T, **Inaba T**, Maki H, Yao A, Kinugawa K, Komuro I.
Heart Vessels. 2016 Jul;31(7):1206-8.

Opening of native aortic valve accomplished after left ventricular assist device implantation in patients with insufficient preoperative Beta-blocker treatment.

Imamura T, Kinugawa K, Nitta D, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Kyo S, Ono M.
Int Heart J. 2015 May 13;56(3):303-8.

Novel scoring system using postoperative cardiopulmonary exercise testing predicts future explantation of left ventricular assist device.

Imamura T, Kinugawa K, Nitta D, Fujino T, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Kyo S, Ono M.
Circ J. 2015;79(3):560-6.

Targeted therapy is required for management of pulmonary arterial hypertension after defect closure in adult patients with atrial septal defect and associated pulmonary arterial hypertension.

Fujino T, Yao A, Hatano M, **Inaba T**, Muraoka H, Minatsuki S, Imamura T, Maki H, Kinugawa K, Ono M, Nagai R, Komuro I.

Int Heart J. 2015;56(1):86-93.

Parasympathetic reinnervation accompanied by improved post-exercise heart rate recovery and quality of life in heart transplant recipients.

Imamura T, Kinugawa K, Okada I, Kato N, Fujino T, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Kyo S, Ono M.

Int Heart J. 2015;56(2):180-5.

Late rejection occurred in recipients who experienced acute cellular rejection within the first year after heart transplantation.

Imamura T, Kinugawa K, Nitta D, Fujino T, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Yao A, Kyo S, Ono M.

Int Heart J. 2015;56(2):174-9.

Platelet-derived growth factor receptor-tyrosine kinase inhibitor, imatinib, is effective for treating pulmonary hypertension induced by pulmonary tumor thrombotic microangiopathy.

Minatsuki S, Miura I, Yao A, Abe H, Muraoka H, Tanaka M, Imamura T, **Inaba T**, Maki H, Hatano M, Kinugawa K, Yao T, Fukayama M, Nagai R, Komuro I.

Int Heart J. 2015;56(2):245-8.

Novel scoring system using cardiopulmonary exercise testing predicts prognosis in heart failure patients receiving guideline-directed medical therapy.

Imamura T, Kinugawa K, Nitta D, **Inaba T**, Maki H, Hatano M, Komuro I.

Circ J. 2015;79(5):1068-75.

Readmission due to driveline infection can be predicted by new score by using serum albumin and body mass index during long-term left ventricular assist device support.

Imamura T, Kinugawa K, Nitta D, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Kyo S, Ono M.

J Artif Organs. 2015 Jun;18(2):120-7.

Is the internal jugular vein or femoral vein a better approach site for endomyocardial biopsy in heart transplant recipients?

Imamura T, Kinugawa K, Nitta D, Fujino T, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Yao A, Kyo S, Ono M.

Int Heart J. 2015;56(1):67-72.

Low blood pressure, low serum cholesterol and anemia predict early necessity of ventricular assist device implantation in patients with advanced heart failure at the time of referral from non-ventricular assist device institutes.

Fujino T, Kinugawa K, Hatano M, Imamura T, Muraoka H, Minatsuki S, **Inaba T**, Maki H, Kinoshita O, Nawata K, Yao A, Ono M, Komuro I.

Circ J. 2014;78(12):2882-9.

Aortic insufficiency in patients with sustained left ventricular systolic dysfunction after axial flow assist device implantation.

Imamura T, Kinugawa K, Fujino T, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Kyo S, Ono M.

Circ J. 2015;79(1):104-11.

Low Blood Pressure, Low Serum Cholesterol and Anemia Predict Early Necessity of Ventricular Assist Device Implantation in Patients With Advanced Heart Failure at the Time of Referral From Non-Ventricular Assist Device Institutes.

Fujino T, Kinugawa K, Hatano M, Imamura T, Muraoka H, Minatsuki S, **Inaba T**, Maki H, Kinoshita O, Nawata K, Yao A, Ono M, Komuro I.

Circ J. 2014 Oct 30.

Lower rotation speed stimulates sympathetic activation during continuous-flow left ventricular assist device treatment.

Imamura T, Kinugawa K, Nitta D, Fujino T, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Kyo S, Ono M.

J Artif Organs. 2015 Mar;18(1):20-6.

Recipients with shorter cardiopulmonary bypass time achieve improvement of parasympathetic reinnervation within 6 months after heart transplantation.

Imamura T, Kinugawa K, Fujino T, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Kyo S, Ono M.

Int Heart J. 2014;55(5):440-4.

Low cardiac output stimulates vasopressin release in patients with stage d heart failure.

Imamura T, Kinugawa K, Hatano M, Fujino T, **Inaba T**, Maki H, Kinoshita O, Nawata K, Kyo S, Ono M,

Komuro I.

Circ J. 2014;78(9):2259-67.

Increased urine aquaporin-2 relative to plasma arginine vasopressin is a novel marker of response to tolvaptan in patients with decompensated heart failure.

Imamura T, Kinugawa K, Fujino T, **Inaba T**, Maki H, Hatano M, Yao A, Komuro I.

Circ J. 2014;78(9):2240-9.

Status 2 patients had poor prognosis without mechanical circulatory support.

Imamura T, Kinugawa K, Hatano M, Fujino T, **Inaba T**, Maki H, Kinoshita O, Amiya E, Nawata K, Yao A, Kyo S, Ono M, Komuro I.

Circ J. 2014;78(6):1396-404.

Differential impacts of achieved heart rate and achieved dose of β -blocker on clinical outcomes in heart failure with and without atrial fibrillation.

Kato N, Kinugawa K, Teruhiko I, Hironori M, Hisataka M, **Inaba T**, Masaru H, Atsushi Y, Nagai R.

Int J Cardiol. 2014 May 1;173(2):331-3.

Cardiac allograft vasculopathy can be distinguished from donor-transmitted coronary atherosclerosis by optical coherence tomography imaging in a heart transplantation recipient: double layered intimal thickness.

Imamura T, Kinugawa K, Murasawa T, Kagami Y, Endo M, Muraoka H, Fujino T, **Inaba T**, Maki H, Hatano M, Kinoshita O, Nawata K, Kyo S, Komuro I, Ono M.

Int Heart J. 2014;55(2):178-80.

Urine sodium excretion after tolvaptan administration is dependent upon baseline serum sodium levels: a possible explanation for the improvement of hyponatremia with scarce chance of hypernatremia by a vasopressin receptor antagonist.

Imamura T, Kinugawa K, Minatsuki S, Muraoka H, Kato N, **Inaba T**, Maki H, Hatano M, Yao A, Komuro I.

Int Heart J. 2014;55(2):131-7.

Late-onset right ventricular failure in patients with preoperative small left ventricle after implantation of continuous flow left ventricular assist device.

Imamura T, Kinugawa K, Kato N, Muraoka H, Fujino T, **Inaba T**, Maki H, Kinoshita O, Hatano M, Kyo S, Ono M.

Circ J. 2014;78(3):625-33.

Preoperative beta-blocker treatment is a key for deciding left ventricular assist device implantation strategy as a bridge to recovery.

Imamura T, Kinugawa K, Hatano M, Fujino T, Muraoka H, Inaba T, Maki H, Kagami Y, Endo M, Kinoshita O, Nawata K, Kyo S, Ono M.

J Artif Organs. 2014 Mar;17(1):23-32.

Tolvaptan can improve clinical course in responders.

Imamura T, Kinugawa K, Minatsuki S, Muraoka H, Kato N, Inaba T, Maki H, Hatano M, Yao A, Komuro I.

Int Heart J. 2013;54(6):377-81.

Successful treatment of hemodynamic compromise caused by antibody-mediated and cellular rejection in a recipient 12 years after heart transplantation.

Imamura T, Kinugawa K, Kato N, Kagami Y, Endo M, Kaneko N, Minatsuki S, Muraoka H, Inaba T, Maki H, Hatano M, Doi K, Yao A, Takazawa Y, Ono M, Kyo S, Komuro I.

Int Heart J. 2013;54(5):328-31.

Everolimus-incorporated immunosuppressant strategy improves renal dysfunction while maintaining low rejection rates after heart transplantation in Japanese patients.

Imamura T, Kinugawa K, Ono M, Kagami Y, Endo M, Minatsuki S, Muraoka H, Kato N, Inaba T, Maki H, Hatano M, Yao A, Kyo S, Komuro I.

Int Heart J. 2013;54(4):222-7.

A case with recovery of response to tolvaptan associated with remission of acute kidney injury and increased urine osmolality.

Imamura T, Kinugawa K, Kato N, Minatsuki S, Muraoka H, Inaba T, Maki H, Shiga T, Hatano M, Hosoya Y, Takahashi M, Yao A, Kyo S, Ono M, Komuro I.

Int Heart J. 2013;54(2):115-8.

Acute pulmonary vasoreactivity test with sildenafil or nitric monoxide before left ventricular assist device implantation.

Imamura T, Kinugawa K, Hatano M, Kato N, Minatsuki S, Muraoka H, Inaba T, Maki H, Kimura M, Kinoshita O, Shiga T, Yao A, Kyo S, Ono M, Komuro I.

J Artif Organs. 2013 Sep;16(3):389-92.

Successful conversion from thiazide to tolvaptan in a patient with stage d heart failure and chronic kidney disease before heart transplantation.

Imamura T, Kinugawa K, Kato N, Minatsuki S, Muraoka H, **Inaba T**, Maki H, Shiga T, Hatano M, Yao A, Kyo S, Ono M, Komuro I.

Int Heart J. 2013;54(1):48-50.

Urine osmolality estimated using urine urea nitrogen, sodium and creatinine can effectively predict response to tolvaptan in decompensated heart failure patients.

Imamura T, Kinugawa K, Minatsuki S, Muraoka H, Kato N, **Inaba T**, Maki H, Shiga T, Hatano M, Yao A, Kyo S, Komuro I.

Circ J. 2013;77(5):1208-13.

Trend of clinical outcome and surrogate markers during titration of β -blocker in heart failure patients with reduced ejection fraction: relevance of achieved heart rate and β -blocker dose.

Kato N, Kinugawa K, Imamura T, Muraoka H, Minatsuki S, **Inaba T**, Maki H, Shiga T, Hatano M, Yao A, Komuro I, Nagai R.

Circ J. 2013;77(4):1001-8.

Correction of hyponatremia by tolvaptan before left ventricular assist device implantation.

Imamura T, Kinugawa K, Shiga T, Kato N, Endo M, **Inaba T**, Maki H, Hatano M, Yao A, Hirata Y, Nishimura T, Kyo S, Ono M, Nagai R.

Int Heart J. 2012;53(6):391-3.

Rapidly progressive cardiac allograft vasculopathy in early onset regressed with everolimus treatment in an adult cardiac recipient.

Ishida J, Kinugawa K, Shiga T, Imamura T, Hatano M, Maki H, **Inaba T**, Yao A, Hirata Y, Nishimura T, Kyo S, Ono M, Nagai R.

Int Heart J. 2012;53(6):388-90.

Novel criteria of urine osmolality effectively predict response to tolvaptan in decompensated heart failure patients--association between non-responders and chronic kidney disease.

Imamura T, Kinugawa K, Shiga T, Kato N, Muraoka H, Minatsuki S, **Inaba T**, Maki H, Hatano M, Yao A, Kyo S, Nagai R.

Circ J. 2013;77(2):397-404.

Bosentan improved persistent pulmonary hypertension in a case after implantation of a left ventricular assist device.

Imamura T, Kinugawa K, Hatano M, Kato N, Minatsuki S, Muraoka H, **Inaba T**, Maki H, Shiga T, Yao A, Kyo S, Ono M, Nagai R.

J Artif Organs. 2013 Mar;16(1):101-4.

Preoperative levels of bilirubin or creatinine adjusted by age can predict their reversibility after implantation of left ventricular assist device.

Imamura T, Kinugawa K, Shiga T, Endo M, Kato N, **Inaba T**, Maki H, Hatano M, Yao A, Nishimura T, Hirata Y, Kyo S, Ono M, Nagai R.

Circ J. 2013;77(1):96-104.

Combination evaluation of preoperative risk indices predicts requirement of biventricular assist device.

Shiga T, Kinugawa K, Imamura T, Kato N, Endo M, **Inaba T**, Maki H, Hatano M, Yao A, Nishimura T, Hirata Y, Kyo S, Ono M, Nagai R.

Circ J. 2012;76(12):2785-91.

How to demonstrate the reversibility of end-organ function before implantation of left ventricular assist device in INTERMACS profile 2 patients?

Imamura T, Kinugawa K, Shiga T, Kato N, Endo M, **Inaba T**, Maki H, Hatano M, Yao A, Hirata Y, Akahane M, Nishimura T, Kyo S, Ono M, Nagai R.

J Artif Organs. 2012 Dec;15(4):395-8.

A case of pancreatic cancer after heart transplantation.

Imamura T, Kinugawa K, Mohri D, Shiga T, Endo M, **Inaba T**, Maki H, Hatano M, Isayama H, Yao A, Hirata Y, Koike K, Nagai R.

Int Heart J. 2012;53(3):205-7.

Successful conversion to everolimus after cytomegalovirus infection in a heart transplant recipient.

Imamura T, Shiga T, Kinugawa K, Kato N, Endo M, **Inaba T**, Maki H, Hatano M, Yao A, Hirata Y, Nagai R.

Int Heart J. 2012;53(3):199-201.

An elevated ratio of early to late diastolic filling velocity recovers after heart transplantation in a time-dependent manner.

Imamura T, Kinugawa K, Shiga T, Endo M, Kato N, **Inaba T**, Maki H, Hatano M, Yao A, Nishimura T, Hirata Y, Kyo S, Ono M, Nagai R.

J Cardiol. 2012 Oct;60(4):295-300.

Novel risk scoring system with preoperative objective parameters gives a good prediction of 1-year mortality in patients with a left ventricular assist device.

Imamura T, Kinugawa K, Shiga T, Endo M, Kato N, **Inaba T**, Maki H, Hatano M, Yao A, Nishimura T, Hirata Y, Kyo S, Ono M, Nagai R.

Circ J. 2012;76(8):1895-903.

Early decision for a left ventricular assist device implantation is necessary for patients with modifier A.

Imamura T, Kinugawa K, Shiga T, Endo M, **Inaba T**, Maki H, Hatano M, Imai Y, Yao A, Hirata Y, Nishimura T, Kyo S, Ono M, Nagai R.

J Artif Organs. 2012 Sep;15(3):301-4.

A case of late-onset right ventricular failure after implantation of a continuous-flow left ventricular assist device.

Imamura T, Kinugawa K, Shiga T, Endo M, **Inaba T**, Maki H, Hatano M, Yao A, Nishimura T, Hirata Y, Kyo S, Ono M, Nagai R.

J Artif Organs. 2012 Jun;15(2):200-3.

Initial and programmed combination therapy with oral drugs for severe idiopathic pulmonary arterial hypertension.

Maki H, Yao A, **Inaba T**, Shiga T, Hatano M, Kinugawa K, Yamashita T, Aizawa T, Nagai R.

Int Heart J. 2011;52(5):323-6.

Intravascular ultrasound, angioscopic and histopathological characterisation of heterogeneous patterns of restenosis after sirolimus-eluting stent implantation: insights into potential "thromborestenosis" phenomenon.

Oikawa Y, Yajima J, Costa MA, Matsuno S, Akabane M, Funada R, **Inaba T**, Nakagawa Y, Nakamura M, Nagashima K, Kirigaya H, Ogasawara K, Sawada H, Aizawa T.

EuroIntervention. 2010 Aug;6(3):380-7.

Clinical outcomes after percutaneous peripheral intervention for chronic total occlusion of superficial femoral arteries: comparison between self-expandable nitinol stent and stainless steel

stent.

Nakagawa Y, Yajima J, Oikawa Y, Ogasawara K, Kirigaya H, Nagashima K, Funada R, Matsuno S, **Inaba T**, Nakamura M, Sawada H, Aizawa T.

J Cardiol. 2009 Jun;53(3):417-21.

The potential of RF backscattered IVUS data and multidetector-row computed tomography images for tissue characterization of human coronary atherosclerotic plaques.

Funada R, Oikawa Y, Yajima J, Kirigaya H, Nagashima K, Ogasawara K, Matsuno S, **Inaba T**, Nakagawa Y, Nakamura M, Kurabayashi M, Aizawa T.

Int J Cardiovasc Imaging. 2009 Jun;25(5):471-8.

Short- or long-term outcomes of coronary artery aneurysms occurring after directional coronary atherectomy.

Oikawa Y, Yajima J, Angiolillo DJ, Akabane M, Funada R, Matsuno S, **Inaba T**, Nakagawa Y, Nakamura M, Sawada H, Aizawa T.

J Invasive Cardiol. 2008 Apr;20(4):159-60.

Introns in protein-coding genes in Archaea.

Watanabe Y, Yokobori S, **Inaba T**, Yamagishi A, Oshima T, Kawarabayasi Y, Kikuchi H, Kita K.

FEBS Lett. 2002 Jan 2;510(1-2):27-30.